Recitation notes on Kneser-Ney

Maria Ryskina

September 7, 2018

1 Notation

- V – corpus vocabulary
- $c(x)$ – count of n-gram x in the corpus
- $N_{1+}(\bullet w) \triangleq |\{u : c(u, w) > 0\}|$ – number of unique bigrams in the corpus ending in w
- $N_{1+}(w\bullet) \triangleq |\{u : c(w, u) > 0\}|$ – number of unique bigrams in the corpus starting with w
- $N_{1+}(\bullet w\bullet) \triangleq |\{(u, v) : c(u, w, v) > 0\}|$ – number of unique trigrams in the corpus with w in the middle
- $\mathbb{1}[\cdot]$ – indicator function

2 Conditional n-gram probabilities

$$P(w|\text{prev}_{k-1}) = \frac{\max(c'(\text{prev}_{k-1}, w) - d, 0)}{\sum_{v \in V} c'(\text{prev}_{k-1}, v)} + \alpha(\text{prev}_{k-1}) P(w|\text{prev}_{k-2}) \tag{1}$$

Highest order (trigram):

$$P(w_3|w_1w_2) = \frac{\max(c'(w_1w_2w_3) - d, 0)}{\sum_{v \in V} c'(w_1w_2v)} + \alpha(w_1w_2) P(w_3|w_2) \tag{2}$$

Lower order (bigram and unigram):

$$P(w_3|w_2) = \frac{\max(c'(w_2w_3) - d, 0)}{\sum_{v \in V} c'(w_2v)} + \alpha(w_2) P(w_3) \tag{3}$$

$$P(w_3) = \frac{c'(w_3)}{\sum_{v \in V} c'(v)} \tag{4}$$

Remembering the definition of $c'(x)$:

- if x is a trigram, $c'(x) = c(x)$ (count of the trigram in the corpus)
- if x is a bigram or a unigram: $c'(x) = N_{1+}(\bullet x)$ (number of unique words preceding x in the corpus)

We substitute it in Equations 2-4:

$$\begin{aligned} P(w_3|w_1w_2) &= \frac{\max(c(w_1w_2w_3) - d, 0)}{\sum_{v \in V} c(w_1w_2v)} + \alpha(w_1w_2)P(w_3|w_2) = \\ &= \frac{\max(c(w_1w_2w_3) - d, 0)}{c(w_1w_2)} + \alpha(w_1w_2)P(w_3|w_2) \end{aligned} \tag{5}$$

$$\begin{aligned} P(w_3|w_2) &= \frac{\max(N_{1+}(\bullet w_2w_3) - d, 0)}{\sum_{v \in V} N_{1+}(\bullet w_2v)} + \alpha(w_2)P(w_3) = \\ &= \frac{\max(N_{1+}(\bullet w_2w_3) - d, 0)}{N_{1+}(\bullet w_2\bullet)} + \alpha(w_2)P(w_3) \end{aligned} \tag{6}$$

$$P(w_3) = \frac{N_{1+}(\bullet w_3)}{\sum_{v \in V} N_{1+}(\bullet v)} = \frac{N_{1+}(\bullet w_3)}{N_{1+}(\bullet\bullet)} \tag{7}$$

Here $N_{1+}(\bullet\bullet)$ is the number of all unique bigrams.

3 Computing α

To compute α, we sum over both sides of Equations 5-6 and use the fact that $\sum_{w \in V} P(w_3 = w|\ldots) = 1$. For the trigram case:

$$\sum_{w \in V} P(w_3 = w|w_1w_2) = \frac{\sum_{w \in V} \max(c(w_1w_2w) - d, 0)}{c(w_1w_2)} + \alpha(w_1w_2) \sum_{w \in V} P(w_3 = w|w_2)$$

$$1 = \frac{\sum_{w \in V} \max(c(w_1w_2w) - d, 0)}{c(w_1w_2)} + \alpha(w_1w_2) \tag{8}$$

Since $0 < d < 1$, we can rewrite this equation as:

$$\begin{aligned} 1 &= \frac{\sum_{w \in V} c(w_1w_2w) - d \cdot \sum_{w \in V} \mathbb{1}[c(w_1w_2w) > 0]}{c(w_1w_2)} + \alpha(w_1w_2) = \\ &= \frac{\sum_{w \in V} c(w_1w_2w) - d \cdot N_{1+}(w_1w_2\bullet)}{c(w_1w_2)} + \alpha(w_1w_2) = \\ &= 1 - \frac{d \cdot N_{1+}(w_1w_2\bullet)}{c(w_1w_2)} + \alpha(w_1w_2) \end{aligned} \tag{9}$$

Finally,

$$\alpha(w_1 w_2) = d \cdot \frac{N_{1+}(w_1 w_2 \bullet)}{c(w_1 w_2)} \tag{10}$$

Now, doing the same for the bigram case:

$$\begin{aligned} 1 &= \frac{\sum_{w \in V} \max(N_{1+}(\bullet w_2 w) - d, 0)}{N_{1+}(\bullet w_2 \bullet)} + \alpha(w_2) = \\ &= \frac{\sum_{w \in V} N_{1+}(\bullet w_2 w) - d \cdot \sum_{w \in V} \mathbb{1}[N_{1+}(\bullet w_2 w) > 0]}{N_{1+}(\bullet w_2 \bullet)} + \alpha(w_2) \end{aligned} \tag{11}$$

Indicator $\mathbb{1}[N_{1+}(\bullet w_2 w) > 0]$ is equal to 1 for every w for which $w_2 w$ occurs in at least one context. That is equivalent to saying bigram $w_2 w$ occurs at least once, so we can replace $\mathbb{1}[N_{1+}(\bullet w_2 w) > 0]$ with $\mathbb{1}[c(w_2 w) > 0]$:

$$1 = 1 - d \cdot \frac{\sum_{w \in V} \mathbb{1}[c(w_2 w) > 0]}{N_{1+}(\bullet w_2 \bullet)} + \alpha(w_2) = 1 - d \cdot \frac{N_{1+}(w_2 \bullet)}{N_{1+}(\bullet w_2 \bullet)} + \alpha(w_2) \tag{12}$$

Finally,

$$\alpha(w_2) = d \cdot \frac{N_{1+}(w_2 \bullet)}{N_{1+}(\bullet w_2 \bullet)} \tag{13}$$

4 Edge cases

- Our derivation until now assumed that $c(w_1 w_2) > 0$, otherwise the denominators turn into 0. If the context $w_1 w_2$ has never occurred before, fully back off to lower order until you get to a context with non-zero count.
- If w_3 is a word that has not been seen before, you can return a zero probability or back off to a uniform model and return $\frac{1}{|V|}$. See Implementation tips for practical advice.

5 Implementation tips

- In your hashmap structures, you might want to store tables for values used for computing α and P in addition to count tables:
  - for every occurring unigram w you would store $N_{1+}(\bullet w)$, $N_{1+}(w \bullet)$, $N_{1+}(\bullet w \bullet)$.
  - for every occurring bigram vw you would store $N_{1+}(vw \bullet)$ and $N_{1+}(\bullet vw)$
- To account for unknown words in translation, you can return a very small constant instead of a zero probablilty in case of a unigram not seen before.